


Appendix A

Laboratory of Gene Expression
University of Aarhus
Department of Molecular and
Structural Biology
Gustav Wieds Vej 10
8000 Aarhus C, Denmark

Biochemistry

FOURTH EDITION



Lubert Stryer

STANFORD UNIVERSITY



W. H. Freeman and Company
New York

Library of Congress Cataloging-in-Publication Data

Stryer, Lubert.
Biochemistry/Lubert Stryer.—4th ed.

Includes index.

ISBN 0-7167-2009-4

1. Biochemistry. I. Title.

QP514.2.S66 1995

574.19'2—dc20

94-22832

©1975, 1981, 1988, 1995 by Lubert Stryer


No part of this book may be reproduced by any mechanical, photographic, or electronic process, or in the form of a phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without written permission from the publisher.

Printed in the United States of America

Second printing 1995, KP

Figure 15-11

Three-dimensional structure of the S1 head of myosin determined by x-ray crystallography at 2.8-Å resolution. The nucleotide-binding site is empty in this crystal form. The heavy chain of S1 consists of a 25-kd N-terminal domain (purple), a 50-kd central domain (blue), and a 20-kd C-terminal domain (red). The actin-binding site and the ATPase site are on opposite sides of the 50-kd domain. Two light chains (RLC, yellow, and ELC, green) are bound to the long C-terminal α helix of the heavy chain. [Courtesy of Dr. Ivan Rayment and Dr. Hazel Holden.]



ments, whereas the actin-binding site is formed by the 50-kd and 20-kd segments. Nearly all known myosins share the catalytic-site sequence



which is similar to the sequences found in the active sites of other ATPases. The two light chains wrap around the 20-kd segment, which forms an 85-Å-long α helix that spans much of S1.

The two S1 heads are joined at a hinge to the tail of myosin, which is a two-stranded coiled coil (Figure 15-12). The formation of this very long rod (1700 Å, or 170 nm) is favored by the absence of proline over a span of more than a thousand residues and by the abundance of leucine, alanine, and glutamate. The two strands are in register, pointing the same way. Their axes are about 10 Å apart, enabling the side chains of the two strands to interact intimately to reinforce the helical structure. Regularities in the amino acid sequence of the tail promote the formation of a coiled-coil rod. The tail consists of repeating seven-residue units (*abcdefg*) in which *a* and *d* are usually hydrophobic (Figure 15-13). Residues *a* and *d* form a zigzag pattern of knobs and holes that interlock with those of the other strand to form a tight-fitting hydrophobic core. In contrast, residues *b*, *c*, and *f*, which are located on the periphery of the coiled coil, tend to be charged.



Figure 15-12
Model of a two-stranded α -helical coiled coil.




Figure 15-13
Schematic diagram of a cross section of an α -helical coiled coil showing the positions of side chains. Residues *a* and *d* (yellow) of each strand pack tightly to form a hydrophobic core. Residues *b*, *c*, and *f* (green) on the periphery tend to be charged.